

流産した女性の非嘆体験と精神的苦痛尺度の考案

著者	島田 啓子
著者別表示	Shimada Keiko
雑誌名	平成15(2003)年度 科学研究費補助金 基盤研究(C) 研究成果報告書概要
巻	2002 2003
ページ	2p.
発行年	2005-04-18
URL	http://doi.org/10.24517/00063681



2003 Fiscal Year Final Research Report Summary

Study on the development of the perinatal grief scale : Women experience a medically-induced miscarriage during the early pregnancy stage.

Research Project

Project/Area Number

14572269

Research Category

Grant-in-Aid for Scientific Research (C)

Allocation Type

Single-year Grants

Section

一般

Research Field

Clinical nursing

Research Institution

KANAZAWA UNIVERSITY

Principal Investigator

SHIMADA Keiko Kanazawa Univ., Faculty of Medicine, Associate professor, 医学部, 助教授 (60115243)

Co-Investigator(Kenkyū-buntansha)

SAKAI Akemi Kanazawa Univ., Faculty of Medicine, Professor, 医学部, 教授 (90115238)
KAMEDA Yukie Kanazawa Univ., Faculty of Medicine, Assistants, 医学部, 助手 (40313671)
IGARASHI Touko Jouetsu Univ., School psychology&counseling, Associate professor, 学校教育学部, 助教授 (90293349)

Project Period (FY)

2002 – 2003

Keywords

Miscarriage / Early pregnancy / Perinatal grief / Scale / Semi-structured interview

Research Abstract

This study focused on the development of the perinatal grief experiences among women who experienced a medically-induced miscarriage during the early pregnancy stage. In 2002, researchers reviewed literatures, and analyzed the perinatal bereavement scale and the perinatal grief scale that were widely employed in the U.S.A. The semi-structure interview was developed and 9 women were interviewed. Buffer or facilitation factors of overcoming the perinatal grief experiences included social support from their partners and/or friends, their hobby and/or work whereas obstacle factors were medical practitioners insensitive behavior.

In 2003, the Japanese perinatal grief inventory scale(J-PGIS) was developed based on these results in order to examine women's perinatal grief experiences. Thirty six women were interviewed with J-PGIS and Zung's depression scale. Cronbach's was .656, and J-PGIS was significantly correlated to the depression scale($r=.646$, $p<.001$). Results suggested the importance of the development of a new care program including a peer support and other medical professionals' involvement. Future study is necessary to obtain more data with ethical considerations in order to establish a more reliable and valid scale.

Research Products (2 results)

AllOther

AllPublications

- [Publications] Shimada Keiko, Igarashi Touko, Kameda Yukie, Sakai Akemi: "Study on the women experience a medically-induced miscarriage during the early pregnancy stage."Journal of Japan Academy of Midwifery. 18・3(predetermine of presentation). (2005) ▼
- [Publications] Shimada Keiko, Igarashi Touko, Kameda Yukie, Sakai Akemi: "Study on the development of the perinatal grief scale with miscarriage."Journal of Japan Academy of Midwifery. 18・3(predetermine of presentation). (2005) ▼

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-14572269/145722692003kenkyu_seika_hokoku_

Published: 2005-04-18